arXiv:1104.0833v1 [math.CV] 5 Apr 2011

# A REMARK ON TWO EXTENSIONS OF THE DISC ALGEBRA AND MERGELYAN'S THEOREM

V. Nestoridis and I. Papadoperakis

## Abstract

We investigate the set of uniform limits of polynomials on any closed Jordan domain with respect to the chordal metric $\chi$ on $\mathbb{C} \cup \{\infty\}$. We conclude that Mergelyan's Theorem may be extended to the case of uniform approximation with respect to $\chi$ on closed Jordan domains. Similar results are obtained if we replace the one point compactification $\mathbb{C}\cup\{\infty\}$ of $\mathbb{C}$ by another compactification of $\mathbb{C}$ homeomorphic to the closed unit disc.

AMS Classification Number: Primary 30J99, secondary 46A99, 30E10.

Key words: Caratheodory theorem, spherical approximation, Mergelyan's Theorem, Disc algebra.

## 1. Spherical approximation

In [2] it has been considered the following generelization $\tilde{A}(D)$ of the disc algebra. $\tilde{A}(D)$ contains the constant function $f(z) \equiv \infty$ for all $z \in \bar{D}$, where $\bar{D}$ is the closed unit disc in $\mathbb{C}$ and all functions $f : D \longrightarrow \mathbb{C}$ holomorphic in the open unit disc $D$, such that, for every $\zeta \in \partial D$ the limit $\lim\limits_{\substack{z\to\zeta \\ z\in D}} f(z)$ exists in $\mathbb{C} \cup \{\infty\}$. It has also been proved ([2]) that $\tilde{A}(D)$ concides with the set of uniform limits with respect to the chordal metric $\chi$ of all polynomials on $\bar{D}$.

Now let $\Omega$ be an open Jordan domain in $\mathbb{C}$ and $\bar{\Omega}$ its closure. Let $\phi : D \longrightarrow \Omega$ be a Riemann map. According to a theorem of Caratheodory ([1]), $\phi$ extends to a homeomorphism $\phi : \bar{D} \longrightarrow \bar{\Omega}$. We consider the set of functions $f \circ \phi^{-1} : \bar{\Omega} \longrightarrow \mathbb{C} \cup \{\infty\}$ for all $f \in \tilde{A}(D)$. It is easily seen that this set coincides with $\tilde{A}(\Omega)$, where $\tilde{A}(\Omega)$ is defined as follows. $\tilde{A}(\Omega)$ contains the function $g(z) \equiv \infty$ on $\bar{\Omega}$ and the functions $g : \bar{\Omega} \longrightarrow \mathbb{C} \cup \{\infty\}$ continous on $\bar{\Omega}$ such that $g(\Omega) \subset \mathbb{C}$ and $g_{|\Omega}$ is holomorphic in $\Omega$.

Theorem 1. Under the above assumptions and notation $\tilde{A}(\Omega)$ coincides with the set of uniform limits with respect to the metric $\chi$ of polynomials on $\bar{\Omega}$.

Proof. Let $P_n$ be a sequence of polynomials and $g : \bar{\Omega} \longrightarrow \mathbb{C} \cup \{\infty\}$ a function such that $\sup\limits_{z \in \bar{\Omega}} \chi(P_n(z), g(z))) \to 0$, as $n \to +\infty$. Then $\sup\limits_{z \in \bar{D}} \chi(P_n \circ \phi(z), g \circ \phi(z))) \to 0$, as $n \to +\infty$. Since $P_n \circ \phi \in A(D)$, there exist polynomials $Q_n$ so that $\sup\limits_{z \in \bar{D}} \chi(P_n \circ \phi(z), Q_n(z)) < \frac{1}{n}$. It follows that $\sup\limits_{z \in \bar{D}} \chi(Q_n(z), g \circ \phi(z)) \to 0$, as $n \to +\infty$. Thus $g \circ \phi \in \tilde{A}(D)$, which implies that $g = (g \circ \phi) \circ \phi^{-1} \in \tilde{A}(\Omega)$.

Conversely, let $g \in \tilde{A}(\Omega)$, then $g = f \circ \phi^{-1}$ for some $f \in \tilde{A}(D)$. Therefore, there exists a sequence of polynomials $P_n$ with $\sup\limits_{z \in \bar{D}} \chi(f(z), P_n(z))) \to 0$, as $n \to +\infty$. It follows that $\sup\limits_{z \in \bar{\Omega}} \chi(g(z), P_n \circ \phi^{-1}(z))) \to 0$, as $n \to +\infty$. Since $P_n \circ \phi^{-1} \in A(\bar{\Omega})$, the classical Mergelyan's Theorem implies that there exist polynomials $Q_n$ satisfying $\sup\limits_{z \in \bar{\Omega}} |P_n \circ \phi^{-1}(z) - Q_n(z)| < \frac{1}{n}$. Since for all $a, b \in \mathbb{C}$ we have $\chi(a, b) \leq |a - b|$, it follows $\sup\limits_{z \in \bar{\Omega}} \chi(P_n \circ \phi^{-1}(z), Q_n(z)) < \frac{1}{n}$. The triangle inequality imples $\sup\limits_{z \in \bar{\Omega}} \chi(Q_n(z), g(z)) \to 0$, as $n \to +\infty$, thus g is the uniform limit with respect to $\chi$ of the squence of polynomials $Q_n$ on $\bar{\Omega}$. This completes the proof.

## 2. Another compactification of $\mathbb{C}$

We identify $\mathbb{C}$ with $D$ by the homeomorphism $\mathbb{C} \ni z \longrightarrow \frac{z}{1+|z|} \in D$. Since $\bar{D}$ is a compactification of $D$, it induces a compactification $\bar{\mathbb{C}} = \mathbb{C} \cup \mathbb{C}^{\infty}$, where $\mathbb{C}^{\infty} = \{\infty \cdot e^{i\theta} : \theta \in [0, 2\pi)\}$. The usual Eucledian distance on $\bar{D}$ iduces a metric $d$ on $\bar{C}$ where $d(z, w) = |\frac{z}{1+|z|} - \frac{w}{1+|w|}|$ for $z, w \in \mathbb{C}$, $d(z, \infty \cdot e^{i\theta}) = |\frac{z}{1+|z|} - e^{i\theta}|$ for $z \in \mathbb{C}$, $\theta \in [0, 2\pi)$ and $d(\infty \cdot e^{i\theta}, \infty \cdot e^{i\phi}) = |e^{i\theta} - e^{i\phi}|$ for $\theta, \phi \in [0, 2\pi)$. In [3] it has been investigated the set of uniform limits with respect to the metric $d$ of the polynomials on $\bar{D}$. This set coincider with the class $\bar{A}(D)$ defined as follows. $\bar{A}(D)$ contains continuous functions $f : \bar{D} \longrightarrow \bar{\mathbb{C}}$ of two types. The finite type are those $f$'s such that $f(D) \subset \mathbb{C}$ and $f_{|D}$ is holomorphic. The infinite type are those $f$'s such that $f(\bar{D}) \subset \mathbb{C}^{\infty}$ and $f(z) = \infty \cdot e^{i\theta(z)}$ where $\theta : \bar{D} \longrightarrow \mathbb{R}$ is continuous on $\bar{D}$ and harmonic on $D$.

Let $\Omega \subset \mathbb{C}$ be an open Jordan domain and $\bar{\Omega}$ its closure. We consider $\phi : D \longrightarrow \Omega$ a Riemann map which it is known that it extends to a homeomorphism $\phi : \bar{D} \longrightarrow \bar{\Omega}$. We consider the set of functions $f \circ \phi^{-1} : \bar{\Omega} \longrightarrow \bar{C}$ for all $f \in \bar{A}(D)$. It is easily seen that this set coincides with the class $\bar{A}(\Omega)$ defined as follows: $\bar{A}(\Omega)$ contains continuous functions $g : \bar{\Omega} \longrightarrow \bar{\mathbb{C}}$ of two types. The finite type is those $g$'s with $g(\Omega) \subset \mathbb{C}$ and $g_{|\Omega}$ holomorphic. The infinite type is those $g$'s with $g(z) \in \mathbb{C}^{\infty}$ for all $z \in \bar{\Omega}$ and $g(z) = \infty \cdot e^{i\theta(z)}$ where $\theta : \bar{\Omega} \longrightarrow \mathbb{R}$ is continuous on $\bar{\Omega}$ and harmonic in $\Omega$.

Theorem 2. Under the above assumptions and notation the class $\bar{A}(\Omega)$ coincides with the set of uniform limits with respect to the metric $d$ of polynomials on $\bar{\Omega}$.

The proof is similar to that of Theorem 1 and is ommitted.

## References

[1] Paul Koosis, Introduction to Hp spaces, London Math. Soc. Lecture Note Series 40, Cambridge University Press, Cambridge, London, N. Y., New Rochelle, Melbourne, Sydney.

[2] V. Nestoridis, An extension of the disc algebra, Arxiv: 1009.5364

[3] V. Nestoridis and N. Papadatos, Another extension of the disc algebra, Arxiv: 1012.3674

[4] W. Rudin, Real and complex Analysis, McGraw-Hill, N.Y., St. Louis, San Fransisco, Toronto, London, Syndney.

V. Nestoridis
Department of Mathematics
University of Athens
Panepistemiopolis
157 84 Athens
Greece
e-mail address: vnestor@math.uoa.gr

I. Papadoperakis
Laboratory of Mathematics
Agricultural University of Athens
118 55 Athens
Greece
e-mail: papadoperakis@aua.gr